

Effect of bacterial isolated *Bacillus subtilus*, *Azospirillum brasilences* and agricultural medium on the growth and flowering of the *Ranunculus asiaticus*

Zahraa Abdul Hadi Jaber *

Kawther hadi Abood

Al-Mussaib Technical College, AL-Furat Al-Awsat Technical University, Iraq.

Email: dr.Kawtherhadi@atu.edu.iq

Abstract : The research was conducted in the greenhouse of the Plant Production Techniques Department, Al-Mussaib Technical College for the autumn season 2019-2020. To study the effect of bacterial isolated *Bacillus subtilus*, *Azospirillum brasilences* and agricultural medium on the growth and flowering of the *Ranunculus asiaticus*, ('Aviv Red' Cultivar) in pots 22 cm in diameter, The research included the effect of two factors, the first factor the bio fertilizers bacterial, with four levels (without the bacterial vaccine, *Bacillus subtilus*, *Azospirillum brasilences*, *Azospirillum brasilences* + *Bacillus subtilus*) and which it were symbolized by (B₀, B₁, B₂, B₃) and the second factor, the agricultural medium that included (River soil + peat moss at a ratio of 2:1 , River soil + 3%, decomposed horse waste ,River soil+ 3% spent mushroom compost(*Agaricus bisporus*) , River soil + 5% decomposed horse waste ,River soil+ 5% spent mushroom compost (*Agaricus bisporus*) and which it were symbolized by (A₀, A₁, A₂, A₃, A₄). factor experiment was conducted according to Completely Randomized Design (C.R.D) and by three replicates and each replicate contains (20 treatments), The averages were compared according to the L.S.D test under the 0.05 probability level .The results of this study can be summarized as follows: -

The treatment with bio fertilizers bacterial resulted in a significant increase in all the studied traits. where the treatment of B₃ (mixture) recorded the highest average in plant height, relative chlorophyll content in leaves, percentage of carbohydrates in leaves, number of flowers, The duration of the flowers on the plant, the number of tuberous roots and the results were as follows: 44.80 cm, 35.92 spad unit , 13.62%, 12.87 flowers.plant⁻¹, 17.03 days, 7.92 roots.plant⁻¹ respectively. The results showed the excelled of the agricultural medium treatments, where the A₃ treatment (River soil + 5% decomposed horse waste) recorded the highest average for most the studied traits, plant height, relative chlorophyll content in the leaves, and, The number of flowers, The duration of the flowers on the plant, the number of tuberous roots and the results were as follows, 44.88 cm, 40.72 Spad unit , 14.23%, 13.38 flowers.Plant⁻¹, 18.83 days, 7.82 roots.plant⁻¹. The results of the bi-interaction between the biofertilizer bacterial and the agricultural medium showed significantly excelled in most the studied traits. where the A₃B₃ treatment excelled in the plant height, the relative chlorophyll content in the leaves, , the number of flowers, the duration of the flower stay on the plant, and the results were as follows: 53.17 cm, 45.23 spad, 17.17 flower.plant⁻¹, 23.11 day. The treatment of the bi-interaction treatment A₂B₃, also excelled by giving the highest average number of tuberous roots 9.11 root.plant⁻¹.

Keyword: bio- fertilizers, *Bacillus subtilus*, decomposed horse waste, *Azospirillum brasilences*, *Ranunculus- asiaticus*

1. INTRODUCTION

Ranunculus asiaticus L, which belongs to Ranunculaceae family , is one of the annual winter flowering bulbs that prefers areas with moderate temperatures .The

Mediterranean region is considered the origin country and it grows well in some areas of Iraq [1].Its economic importance is due to the fact that its flowers are suitable for commercial picking, As

flowers are used in flower bouquets, either singly or after coordination with other flowers. Some types of *Ranunculus* are used in alternative medicine as an anti-rheumatism, fever, and redness of the skin, and this comes through the presence of the substances Anemonin and Protoanemonin, which are used as a treatment for many diseases [2,3]. The use of biofertilizers (bacterial) is necessary to limit the use of chemical fertilizers. Among the bacterial fertilizers is the *Bacillus subtilis* bacteria, which is one of the genera that dissolves phosphates through their secretion of organic acids such as acetic, lactic, formic [4]. Likewise, *Azospirillum* bacteria, which affect the plant's nutrition with nitrogen, fix atmospheric nitrogen, convert it into ammonium ion and nitrate, and increase its absorption by the plant [5]. Organic fertilizers of (decomposed horse waste, mushrooms residues) are an important and essential source of the macro and micro elements needed by the plant, in addition to improving the physical, chemical and biological properties of the soil through the breakdown of heavy soil particles, and improving its aeration also contributes to increasing biological activity within the area of root proliferation because it contains some beneficial microbes and stimulates bioprocesses [6]. The sources of organic matter vary from plant sources that come from plant roots and fallen leaves on the surface of the soil, which go through stages of biological decomposition by the action of microorganisms to plants grown to fluctuate in the soil, such as green compost plants and plant wastes that are added to the soil to increase its productivity and improve its qualities and animal sources come as a result of the effectiveness of reviving the soil and its cells and tissues after their death, as well as human and animal wastes added to the soil [7] when using biological fertilizer, the *Ranunculus* plant obtained an

increase in the number of flowers and the diameter of tuberous roots. and due to the importance of expanding the production of ornamental flowers and bulbs, including the *Ranunculus* plant, in Iraq. Therefore, this research aims to study the type of agricultural medium and added different bacterial types of bio fertilizers in the growth and flowering of the *Ranunculus* plant.

2. MATERIALS AND METHODS

The research was conducted in the greenhouse of the Plant Production Techniques Department, Al-Mussaib Technical College for the autumn season 2019-2020, to study the effect of bacterial isolated *Bacillus subtilis*, *Azospirillum brasiliense* and agricultural medium on the growth and flowering of the *Ranunculus* ('Aviv Red' Cultivar), produced by Flower Holland. where the bulbs were soaked in water for 8 hours, and they were planted in plastic pots of 22 cm in diameter, where the experiment included two factors, The first factor includes adding two types of bio fertilizers bacterial loaded on peat moss and produced in the laboratories of the Agricultural Research Department of the Ministry of Science and Technology, namely *Bacillus subtilis* and *Azospirillum brasiliense*. It was added, according to the experiment treatments of (10 g. pot⁻¹). It was taken into account when cultivating that the bacterial fertilizers were in contact with the bulbs by placing them below and above the bulb [8]. Included on four levels are (without the bacterial vaccine, *Bacillus subtilis*, *Azospirillum brasiliense*, *Azospirillum brasiliense* + *Bacillus subtilis*) and which it were symbolized by (B₀, B₁, B₂, B₃). As for the second factor, the agricultural medium that included (River soil + peat moss, River soil + 3%, decomposed horse waste, River soil + 3% spent mushroom compost (*Agaricus bisporus*), River soil + 5% decomposed horse waste, River soil + 5% spent mushroom compost (*Agaricus bisporus*) and which it were symbolized by (A₀, A₁, A₂, A₃, A₄). Horse waste was collected from the stables of the Lebanese Equestrian Club in the Najaf provain.

Experiment was applied according to (Completely Randomized Design (C.R.D) and by three replicates. The averages were compared according to the L.S.D test under the 0.05 probability level[9]. The data were analyzed using the ready-made statistical program Genstat. Samples were taken from agricultural

medium (mushroom residues, decomposed horse waste) and river soil for analysis in the Central Laboratory for Graduate Studies, University of Baghdad, College of Agricultural Engineering Sciences (Table 1 and 2) before distributing it to the treatments.

Table 1. Chemical and physical properties for horse waste and mushroom residues

Traits	Units	mushroom residues	horse waste
Electrical conductivity (EC)	d.S.m ⁻¹	4.4	4.16
PH	---	7.4	7.28
Organic matter	g.Kg ⁻¹	25.40	28.45
The ratio of carbon to nitrogen	---	16.14	15.40
Total nitrogen	g.Kg ⁻¹	8.75	6.56
phosphorous availability	g.Kg ⁻¹	0.17	0.56
Potassium availability	g.Kg ⁻¹	0.04	0.11

*It was analyzed in the Central Laboratories for Graduate Studies, University of Baghdad, College of Agriculture

Table 2.the chemical and physical characteristics of the soil under study

Unites	Values	Measured chemical and physical properties
	7.43	PH
g.Kg ⁻¹	8.31	Organic matter
dsm ⁻¹	4.21	Electrical conductivity (EC)
mg.Kg ⁻¹ soil	28	Nitrogen
	4.14	Phosphorous
	287.12	Potassium
soil separates		
g.Kg ⁻¹ soil	718	Sand
	195	Silt
	87	Clay
loamy sand		Texture

2.1. Studied traits

2.1.1. Plant height (cm)

2.1.2. Relative chlorophyll content in leaves (SPAD unit)

2.1.3. Number of flowers per plant (flower.plant⁻¹)

According to the number of opening flowers for each plant and for all plants of the experimental unit, then the average was calculated.

2.1.4. The duration of the flowers on the plant (day)

Calculated from when each flowering on the plant until the petals begin to wither.

2.1.5. The number of tuberous roots.plant⁻¹

The average of tuberous roots growing for all experimental unit plants was extracted and then averages were calculated for each treatment.

The results in Table (3) indicate that bio fertilizers had a significant effect on increasing plant height, where the mixture treatment B₃ (*Azospirillum brasilences* + *Bacillus subtilus*) was significantly exceeded and gave the highest plant height of 44.80 cm. It was followed by the treatment of the bio fertilizer B₂ (*Azospirillum brasilences*), which gave a height amounted to (41.43) cm, while the control treatment B₀ gave the lowest height of 32.93 cm. The use of agricultural medium had a significant effect on plant height, where the treatment of A₃ (River soil + 5% decomposed horse waste) excelled and gave the highest averages of (44.88) cm. This was followed by the A₂ treatment that gave a height that amounted to (41.88) cm, excelled on the treatment A₄, which gave the lowest height of (33.42) cm. The bi-interaction between bio fertilizers and the agricultural medium achieved a significant increase in the height of the *Ranunculus* plant. The A₃B₃ treatment recorded the highest average of (53.17) cm and thus excelled on all the treatments, while the A₄B₀ treatment recorded the lowest average of plant height of (28.67) cm.

3. RESULTS AND DISCUSSION

3.1. Plant height (cm)

Table 3. The effect of biofertilizer bacterial and the agriculture medium and their interaction on plant height (cm) of *Ranunculus asiaticus*.

The effect of agriculture medium (A)	biofertilizer bacterial				agriculture medium)A(
	Mixture (B3)	<i>Azospirillum brasilences</i> (B2)	<i>Bacillus subtilus</i> (B1)	without the bacterial vaccine B0)(
34.83	39.00	36.33	34.00	30.00	River soil + peat moss (A0)
39.42	45.00	42.33	37.67	32.67	River soil + 3%, decomposed horse waste(A1)
41.88	48.33	44.83	39.17	35.17	River soil+ 3% spent mushroom compost (A2)
44.88	53.17	47.17	41.00	38.17	River soil + 5% decomposed horse waste(A3)
33.42	38.50	36.50	30.00	28.67	River soil+ 5% spent mushroom compost (A4)
	44.80	41.43	36.37	32.93	The effect of biofertilizer bacterial (B)
1.40				A	L.S.D 0.05
1.25				B	
2.81				A*B	

3.2. Relative chlorophyll content in leaves (SPAD unit)

The results in Table (4) showed that the addition of the bio fertilizer led to a significant increase in the relative chlorophyll content, where the treatment B₃ significantly excelled and gave 35.92 Spad , while the control treatment B₀ gave the lowest values of (27.89) Spad. It is also noticed from the results of the same table that the agricultural medium had a significant increase in recorded the highest average, reaching 45.23 Spad.

the relative chlorophyll content in the leaves. The treatment A₃ significantly excelled on the rest of the treatments and gave the highest average of (40.72) Spad. While the A₄ treatment gave the lowest average of (25.44) Spad. The bi-interaction treatment differed significantly between the bio fertilizers and the agricultural medium and all the treatments excelled on control treatment A₄B₀, which gave the lowest chlorophyll content of (22.44) Spad, while the interaction treatment A₃B₃

Table 4. The effect of biofertilizer bacterial and the agriculture medium and their interaction on Relative chlorophyll content in leaves (SPAD unit) of *Ranunculus asiaticus*.

The effect of agriculture medium (A)	biofertilizer bacterial				agriculture medium)A(
	Mixture (B3)	<i>Azospirillum brasilences</i> (B2)	<i>Bacillus subtilus</i> (B1)	without the bacterial vaccine B0)(
25.76	29.49	25.62	24.89	23.04	River soil + peat moss (A0)
29.86	34.53	30.69	28.78	25.45	River soil + 3%, decomposed horse waste(A1)
36.94	40.96	38.87	35.64	32.27	River soil+ 3% spent mushroom compost (A2)
40.72	45.23	42.37	39.03	36.24	River soil + 5% decomposed horse waste(A3)
25.44	29.40	26.33	23.60	22.44	River soil+ 5% spent mushroom compost (A4)
	35.92	32.77	30.39	27.89	The effect of biofertilizer bacterial (B)
4.57				A	L.S.D 0.05
4.08				B	
9.14				A*B	

3.3. Number of flowers (flower.Plant⁻¹)

The results in Table (5) show that the bio fertilizers achieved a significant increase in the number of flowers formed on each plant. The mixture treatment B₃ (*Bacillus subtilus* + *Azospirillum brasilences*) was significantly excelled and gave it the highest number of flowers, which reached 12.87 flower.Plant⁻¹ compared to the control treatment B₀ , which

gave the lowest number of flowers, which reached 7.97 flower.Plant⁻¹. From the results in the same table, we notice the significant effect of the agricultural medium on the number of flowers formed on the plants. where the A₃ treatment (River soil + 5% decomposed horse waste) significantly excelled on the rest of the medium(A₀, A₁, A₂) by recorded the highest number of flowers formed, which reached 13.38 flower.Plant⁻¹ and thus excelled on the control

treatment A₀ (River soil + peat moss), which recorded the lowest numbers, which are 7.50 flower.Plant⁻¹.Where, it showed the bi- interaction between the bacterial fertilizer B₃ (mixture) and the culture medium A₃ (River soil + 5% decomposed horse waste). It achieved the highest average of 17.17 flower.Plant⁻¹, compared to the control treatment A₀B₁ in which the number of flowers formed on plants decreased and reached 6.83 flower.Plant⁻¹.

Table 5. The effect of biofertilizer bacterial and the agriculture medium and their interaction on Number of flowers (flower. Plant⁻¹)of the *Ranunculus asiaticus*.

The effect of agriculture medium (A)	biofertilizer bacterial				agriculture medium)A(
	Mixture (B3)	<i>Azospirillum brasilences</i> (B2)	<i>Bacillus subtilus</i> (B1)	without the bacterial vaccine B0)(
7.50	8.50	7.67	6.83	7.00	River soil + peat moss (A0)
10.38	13.00	9.17	12.00	7.33	River soil + 3%, decomposed horse waste(A1)
10.88	14.83	11.50	9.50	7.67	River soil+ 3% spent mushroom compost (A2)
13.38	17.17	15.33	13.50	7.50	River soil + 5% decomposed horse waste(A3)
10.25	10.83	10.83	9.00	10.33	River soil+ 5% spent mushroom compost (A4)
	12.87	10.90	10.17	7.97	The effect of biofertilizer bacterial (B)
1.51					A
1.35					B
3.02					A*B
					L.S.D 0.05

3.4. The duration of the flowers on the plant(day) The data in Table (6) refer to the role of biological fertilizers in prolonging the duration of the *Ranunculus* flowers on the plant .The treatment of mixture B₃ significantly excelled and gave the highest average of (17.03) days, excelled on the control treatment B₀ that gave (11.88) days. The results of the table also show that the treatment of agricultural medium had an effect on prolonging the duration of flowers on the plant. where the A₃ treatment (River soil + 5%decomposed horse waste) significantly

excelled and gave (18.83) days, While the duration of flowers on the plant decreased to its lowest average in the treatment of A₄ (River soil + spent mushroom compost (*Agaricus bisporus*)5%), amounting to (10.77) days. As it can be noted from the data of the bi- interaction between biofertilizers and agricultural medium, the A₃B₃ interaction treatment was significantly excelled on the rest of the treatments and gave (23.11) days. Where the flowering duration decreased when the interaction treatment A₄B₀ gave (9.19) days.

Table 6. The effect of biofertilizer bacterial and the agriculture medium and their interaction on The duration of the flowers on the plant (day)of the *Ranunculus asiaticus*

The effect of agriculture medium (A)	biofertilizer bacterial				agriculture medium)A(
	Mixture (B3)	<i>Azospirillum brasilences</i> (B2)	<i>Bacillus subtilus</i> (B1)	without the bacterial vaccine B0)(
12.45	15.45	13.11	10.92	10.33	River soil + peat moss (A0)
14.12	16.66	14.96	13.53	11.33	River soil + 3%, decomposed horse waste(A1)
14.72	17.13	14.44	14.33	12.99	River soil+ 3 spent mushroom compost (A2)
18.83	23.11	19.89	16.78	15.55	River soil + 5% decomposed horse waste(A3)
10.77	12.78	11.36	9.77	9.19	River soil+ 5% spent mushroom compost (A4)
	17.03	14.75	13.06	11.88	The effect of biofertilizer bacterial (B)
1.78					A
1.59					B
3.57					A*B
					L.S.D 0.05

Table 7. The effect of biofertilizer bacterial and the agriculture medium and their interaction on Number of tuberous roots . plant⁻¹ of *Ranunculus asiaticus*.

The effect of agriculture medium (A)	biofertilizer bacterial				agriculture medium)A(
	Mixture (B3)	<i>Azospirillum brasilences</i> (B2)	<i>Bacillus subtilus</i> (B1)	without the bacterial vaccine B0)(
5.86	6.78	6.55	5.44	4.66	River soil + peat moss (A0)
7.05	7.44	7.44	7.67	5.67	River soil + 3%, decomposed horse waste(A1)
7.57	9.11	8.39	7.33	5.44	River soil+ 3% spent mushroom compost (A2)
7.82	8.55	8.39	7.77	6.55	River soil + 5% decomposed horse waste(A3)
6.68	7.72	6.45	6.68	5.88	River soil+ 5% spent mushroom compost (A4)
	7.92	7.44	6.98	5.64	The effect of biofertilizer bacterial (B)
1.14					A
1.02					B
2.28					A*B
					L.S.D 0.05

Biofertilizers play an important and effective role in increasing the Studied traits. showed that the treatment of B₃ (*Bacillus subtilis* + *Azospirillum brasilense*) was significantly excelled in traits of plant height , chlorophyll content, number of flowers, duration of flowering and number of tuberous roots . This excelled is due to the role of biofertilizers in improving the root growth of the plant by encouraging the formation of root hairs and increasing the surface of the root system, which leads to an increase in the absorption of water and nutrients and thus improves the vegetative growth of the plant[10,11] as well as the role of *Bacillus* bacteria on Production of plant growth-regulating hormones [12] ,The results of testing the hormonal content in the agriculture medium[13] demonstrated its ability to produce hormone-like substances such as Auxin, gibberellin, and Zeatin that help increase the average of cell division, expansion and elongation, and an increase in the level of metabolism within cells, and this is reflected in plant growth[14], which increases plant height. Bio-fertilizers contribute to increasing the absorption of nutrients, which play an important role in increasing the size and number of leaf cells. In addition to the role of these organisms in the secretion of growth regulators, including gibberellins, which perform their role in building carbohydrates and proteins and increase their accumulation in the plant[15] and that the increase in flowering traits is due to perhaps the reason for the excelled that these types of bacteria belong to the group of growth-stimulating bacteria (PGPR) that increases the concentrations of nutrients due to their role in dissolving phosphate compounds and fixing atmospheric nitrogen and reducing the degree of soil reaction and thus increasing the release of nutrients, including potassium, or by secreting them many growth regulators and organic acids that work to increase the concentration of elements in the soil and thus reflect positively on plant growth[16] ,The results also showed that bio fertilizers had a significant effect in increasing the number of tuberous roots (Tables 8). This is due to its role in improving the biological and chemical properties of the soil by releasing greater amounts of nutrients available

for absorption by the roots, expanding the spread of the roots and absorbing water and nutrients, which leads to an increase in the efficiency of the photosynthesis process [17] .This result agrees with [10] in its experiment on the *Ranunculus* plant when using *Bacillus subtilis* bacteria and with[18] on *gladiolus* when using *Azospirillum brasilense*. And[19] on the *Catharanthus roseus* plant .From the results in Tables (3-8), it is noted that the agricultural medium A₃ (River soil + 5% decomposed horse waste) A₂ (River soil + 3% spent mushroom compost (*Agaricus bisporus*) on A₀ in most of the characteristics of *Ranunculus*. This may be due to the content of the medium with a high percentage of organic matter and nutrients N, P, K compared to the medium A₀ (River soil + Peat moss)., this works to improve soil fertility, provide nutrients, and increase its availability and absorption by the plant, which affects various biological activities such as cell division and elongation, building proteins and nucleic acids, and thus forming good vegetative growth[20] . This result agrees with [21] on *Antirrhinum majus* and with [22] on *Catharanthus roseus* L plant when using spent mushroom compost, ,[23] when using horse waste on pomegranate trees and with [24] on the plant Chinese Cabbage.

4. Conclusions

The mixture of B₃ bacterial Biofertilizer (*Bacillus subtilis* + *Azospirillum brasilense*) achieved the best results by giving it the highest average of vegetative and flowering traits, vase life and the yield of tuberous roots. The agricultural medium A₂ (River soil+ 3% spent mushroom compost (*Agaricus bisporus*) and A₃ River soil + 5% decomposed horse waste)had a significant effect on the trait of vegetative growth, flowering, vase life and the yield of tuberous roots, but the highest average was achieved by the agricultural medium A₃ for all the studied traits.

REFERENCES

- [1]Badr, Mustafa and Mahmoud, Khattab, Tariq, Al-Qai'i, Muhammad Yaqut, Mustafa Raslan, Mustafa Badr, Muhammad Heikal and Alamuddin Noah. (2010) Flowers, ornamental plants, design and landscaping,

- ninth edition, Ma'arif facility in Alexandria.
- [2] Aslam, M.S, Ijaz, A.S. (2012). The genus *ranunculus*: a phytochemical and ethno pharmacological review. International Journal of Pharmacy and Pharmaceutical Science. 4(5): 15–22.
- [3] Bernstein, N., Ioffe, M., Luria, G., Bruner, M., Nishri, Y., Philosoph-Hadas, S., .and Matan, E. (2011). Effects of K and N Nutrition on Function and Production of *Ranunculus asiaticus*. Pedosphere, 21(3), 288-301.
- [4] Al-Serafy, Zakaria Massad. (2010). Fertilizer Chemistry, Faculty of Agriculture, Mansoura University, First Edition, Dar Al Kotob for Publishing and Distribution.
- [5] Vessey,JK.2003.Plant growth promoting rhizobacteria asbiofertilizer .plant and Soil 255,p:571-586.
- [6] AlcaAntara ,B; R. Mary;B. Almudena and L.Francisco(2016).Liquid Organic Fertilizers For Sustainable Agriculture :Nutrient Up.
- [7] Al-Hasnawi, H. A., Hussein, J. K., & Khaleel, T. H. (2019). Effect of growth regulators and preservative solution on vase life and water relation of gladiolus hybrida l. after cut flowers. The Iraqi Journal of Agricultural Science, 50, 182-191.
- [8] Matysiak, B. and Falkowski, G. 2010. Response of three ornamental plants species to inoculation with *Arbuscular mycorrhiza* fungi depending on compost addition to peat substrate and the rate of controlled release fertilizer. J. Fruit and Ornamen. P. Res.,18(2): 321-333
- [9] Granatsteir, D. (2004). Center for sustaining Agriculture and Natural Resources. Washington State University, Wentchee, Wa.USA.
- [10] Al-Sharaa, Marwa Ali Abdul-Jabbar (2020). The effect of biological fertilizers and humic acid on the growth of *Ranunculus asiaticus*. Master thesis. faculty of Agriculture . Al-Qasim Green University. Iraq.
- [11]Khalaf, Zahraa Abdel Sattar. 2013. Effect of bio pollen and organic fertilizer on growth and storage of clove flowers. Master Thesis . faculty of Agriculture . Baghdad University . Iraq.
- [12] Bhat, M.A.; R.Rasool and S.Ramzan.(2019).Plant Growth Promoting Rhizobacteria (PGPR) for Sustainable and Eco-Friendly Agriculture. Acta Scientific Agriculture 3(1):23-25.
- [13] Yadav, K; K.and S. Sarkar. (2019).Biofertilizers ,Impact on soil Fertility and Crop Productivity under Sustainable Agricultur.Environment and Ecology 37(1):89-93.
- [14] GnamamanickS.S.(2006).Plant Associated Bacteria.Springer,NetherlansPP.155-194.
- [15] Swain, M .R.; Naskar,K.S. and Ramesh,C.R.(2007).Indole-3-acetic acid production and effect on sprouting of yam (*Dioscorea rotundata* L.)minisetts by *Bacillus subtilis* isolated form culturable cow dung microflora .Polish Journal of Microbiology,56(2):103-110.
- [16] Hobbie , E.A .2006.Carbon allocation to ectomycorrhiza fungi correlates with total belowground allocation in culture studies. Ecology 87,P:563-569.
- [17] Alkhalil, Sherine Muzaffar Ali, 2011. Tyr Integration between mineral, organic and biological fertilization in tomato yield (*Lycopersicon esculentum* Mill) in greenhouses. Master thesis, Department of Soil Science and Water Resources - University of Baghdad - College of Agriculture
- [18] Contreras-Angulo,J.R.;T.M.Mata;P.Sara and P.Cuellar-Bermudez (2019). Symbiotic Co-Culture of *Scenedesmus* sp. And *Azospirillum brasilense* on N-Deficient Media with Biomass Production for Biofuels.Sustainability ,11(7):1-16. 27(2). 201-205.
- [19] Qasim, M.; A. Younis; Z. A. Zahir; A. Riaz; H. Raza and Tariq, U. 2014. Microbial Inoculation Increases The Nutrient Uptake Efficiency For Quality Production Of *Gladiolus grandifloras*. Pakistan J. Agri. Sci., 51(4): 875–880.
- [20] Sumaysim, Enas Abdul-Raouf Ammar Rabie, Muslim Abd Ali (2013): The effect of bacilloptrazole and the biomodulation

- Bacillus subtilis on the growth of the eye of the bison plant (*Catharanthus roseus* L). Faculty of Agriculture, University of Kufa.
- [21] Al-Sawaf, Muhammad Dawood Salim and Alwan, Nibras Ihsan Abdul-Jabbar (2010). The effect of some agricultural treatment on the growth, flowering and yield of the germs and creams of two varieties of the freesia plant, *Freesia hybrida*. Al-Rafidain Agriculture Journal, Volume 38, Annex 1, pp. 73-78.
- [22] Abdallateef, S. (2015) Influence of chelated Calcium and growing media on the growth and flowering of *Antirrhinum majus*. The Iraqi Journal of Agriculture Sciences, 46(5):813-818
- [23]Khattak, A.; I. Ahmad; N. UlAmi; F. Wahi and Rahman, H. 2011. Effects of different amended organic media on the growth and development of *Vinca rosea* 'Victory'. Sarhad J. Agric., 27(2): 201-205.
- [24] E.laczi. A , Apahidean, E. Luca, A. Dumitra°, P. Boancã.(2016) .Headed Chinese cabbage growth and yield influenced by different manure types in organic farming system.